




Responses of Stomatal Characteristics to Environmental Factors in *Fabaceae* Trees of the Urban Forest in Maros Regency



Received: 13 November 2024

Accepted: 03 March 2025

Published: 14 March 2025

Roslina Yahya¹ , Hadija Sukri¹ , Nirawati¹ ,
Ira Nurhayati Djarot² 

AFFILIATIONS

1. University of Muslim Maros
2. National Innovation Research Agency, South Tangerang

 CC BY 4.0

The response of plants to environmental conditions is reflected in their stomatal characteristics, including stomatal type, number, and density. This novel research examines the influence of environmental factors on the stomatal characteristics of *Fabaceae* tree leaves in the Urban Forest of Maros Regency. The study explores the relationship between stomatal traits and environmental variables such as light intensity and air humidity. Data analysis employed Pearson correlation to assess how these variables affect stomatal number and density. The results reveal variations in stomatal types among *Fabaceae* species: *Acacia* exhibits *diacytic* stomata, Dadap Merah (*Erythrina crista-galli* L) has *paracytic* stomata, Trembesi shows *normocytic* stomata, and Sengon (*Paraserianthes falcataria* L) displays *cyclocytic* stomata. The study also finds that stomatal number and density are higher in the eastern region compared to the western region. Sengon exhibits the highest stomatal number and density among the species studied, whereas Dadap Merah has the lowest. Correlation analysis indicates that only the External Light Intensity of the Stand (ELIS) significantly affects stomatal number and density at the 0.05 (5%) level. Air humidity does not show a significant impact. These findings highlight the role of light intensity in shaping stomatal characteristics in *Fabaceae* trees within urban forests, contributing to a deeper understanding of plant-environment interactions and their ecological implications.

Keywords: Air humidity, *Fabaceae* tree species, Green Open Space, Stomatal characteristics, Light intensity, Urban Forest conservation

1. Introduction

The existence of urban forests is a significant factor influencing social and ecological sustainability in cities (Djahangard et al., 2024). Urban forests enhance the well-being of urban residents by providing ecosystem services such as air purification, temperature regulation, and noise reduction (Nowak et al., 2014). Trees naturally absorb carbon dioxide, produce oxygen, and mitigate heat, leading to cooler ambient air temperatures, while also serving as noise reducers (Handayani et al., 2024). Research has

demonstrated that urban forests can substantially alleviate urban heat island effects and improve air quality by filtering pollutants (Escobedo et al., 2011). The recommended proportion of green open space in urban areas is at least 30%, according to various urban planning guidelines aimed at ensuring environmental sustainability and public health (WHO, 2016). The area of green open space in Maros Regency ranges from six to seven hectares, with 0.5 hectares designated as an Urban Forest. In the Maros Regency Urban Forest, various types of trees exist,

playing an essential role in balancing air quality against activities occurring in urban areas. The diversity of tree species in urban forests bolsters resilience to pests and diseases, thereby ensuring long-term sustainability (Ordóñez & Duinker, 2013). The function of urban forests in Maros Regency plays an important role in maintaining the balance of the urban ecosystem. And one of the plant families commonly found in urban forests is *Fabaceae*, which is known for its high adaptability to various environmental conditions (Firmansyah et al., 2022). Stomata on *Fabaceae* leaves play a key role in the gas exchange process, which affects the efficiency of photosynthesis (Demars et al., 2024). Various environmental factors, such as temperature, humidity, and light intensity, contribute to the formation and distribution of stomata. (Merdekawati, 2015) By understanding the characteristics of stomata on the leaves of *Fabaceae* trees, we can gain deeper insights into the adaptation of plants to the often-changing urban environment. (Suwandi, 2021) stated that green open spaces can balance CO₂ absorption and O₂ provision. Trees located in urban forest areas greatly affect the air around the urban forest, where trees have the ability to photosynthesize and in the photosynthesis process require stomata and chlorophyll (Juslina, 2022). The characteristics of stomata were influenced by the environmental conditions where the plants grow (Sinay & Lesilolo, 2020). The number and density of stomata show the response of plants to the environmental conditions where they grow (Merdekawati, 2015). The density and number of stomata were influenced by environmental conditions, where the density of stomata increases as light intensity increases to maximize carbon dioxide absorption (Budiono et al., 2016). One significant environmental factor is light intensity. Increased light intensity stimulates the process of photosynthesis, which in turn increases the plants' need for carbon dioxide (Chauhan et al., 2023). To meet this need, plants adapt by opening more stomata and increasing their density. Research shows that plants exposed to more light tend to have more and broader stomata, which allows for more efficient gas exchange.

Besides light intensity, air humidity also plays a vital role in the regulation of stomata (Cambaba & Kasi, 2022). In high humidity conditions, plants can open their stomata wider to enhance photosynthesis without worrying about losing too much water. However, in low humidity conditions, stomata will close to reduce evaporation, although this can limit photosynthesis capacity. (Raharjo et al., 2015). Therefore, plants must adapt to changing environmental conditions to balance photosynthesis needs and water conservation.

The varying density and number of stomata among plant species indicate different environmental adaptations (Clark et al., 2022). Stomatal density and distribution are influenced by genetic factors as well as environmental conditions such as light intensity, humidity, and CO₂ concentration (Casson & Gray, 2008). In general, plants that grow in drier habitats have a higher stomatal density in response

to water stress (Haworth et al., 2023; Sulistiyowati et al., 2021). This adaptation allows plants to optimize water use efficiency and maintain photosynthetic activity under limited water availability (Franks & Farquhar, 2007). Additionally, studies have shown that plants in arid environments often exhibit smaller stomata, which further enhances their ability to regulate water loss (Hetherington & Woodward, 2003). These adaptations highlight the critical role of stomata in plant survival and productivity under varying environmental conditions (Lawson & Blatt, 2014). The research aims to determine the influence and how environmental factors affect the stomatal characteristics of *Fabaceae* tree leaves in the Urban Forest of Maros Regency.

2. Method

1) Research Area

This research was conducted in March-August 2024 in the Urban Forest (Green Open Space) of Maros Regency (figure 1), The study was conducted in the Maros Regency Urban Forest, located in South Sulawesi, Indonesia. The urban forest covers an area of approximately 0.5 hectares and is home to a variety of tree species, including several members of the *Fabaceae* family. Analysis of research samples were carried out in the laboratory of the Faculty of Agriculture, Hasanuddin University.




Figure 1. Research location in Maros in the Urban Forest (Green Open Space) of Maros Regency

2) Tools and Materials

The study used tools and materials consisting of Lux Meter, SPAD-502 Plus chlorophyll meter, Binocular Microscope, preparation glass, preparation box (slide box), razor blade (cutter), apparent isolation, pole, digital camera, pen, tally sheet. The materials used in the study consisted of: batteries, clear cuttings, and four types of *Fabaceae* plants, namely: Acacia tree leaves (*Acacia auriculiformis*), Dadap Merah plant leaves (*Erythrina cristagalli* L), Sengon tree leaves (*Paraserianthes falcataria* L), Trembesi tree leaves (*Samanea saman* (Jacq.) Merr.)

3) Data Collection Methods and Instruments

The data collected consists of two types. First, primary data is data obtained directly from observations, measurements, calculations, and data

analysis directly in the field related to the relationship of stomatal characteristics to environmental factors on the leaves of *Fabaceae* trees in the Urban Forest of Maros Regency. The second data is secondary data Secondary data is data obtained from literature studies by searching literature through numbers, scientific articles, books to support research is secondary data in this study.

4) Sampling and Data Collection

Leaf samples were collected from four *Fabaceae* species: *Acacia auriculiformis*, *Erythrina cristagalli* L., *Paraserianthes falcataria* L., and *Samanea saman* (Jacq.) Merr. For each species, five replicates were taken to ensure representative sampling (table 1). Light intensity and air humidity were measured using a Lux Meter and a hygrometer, respectively.

Table 1. Population of *Fabaceae* trees in Maros Regency Urban Forest

| No | <i>Fabaceae species</i> | Population | Sampel | repeatability |
|--------------|--|------------|----------|---------------|
| 1. | Akasia (<i>Acacia auriculiformis</i>) | 1 | 1 | 5 |
| 2. | Dadap merah (<i>Erythrina cristagalli</i> L) | 2 | 1 | 5 |
| 3. | Sengon (<i>Paraserianthes falcataria</i> L) | 2 | 1 | 5 |
| 4. | Trembesi (<i>Samanea saman</i> (Jacq.) Merr.) | 1 | 1 | 5 |
| Total | | 7 | 4 | 20 |

Fabaceae trees in the Maros Regency urban forest have populations that can be seen in table 2, with a total population of *Fabaceae* as many as seven populations with acacia and trambesi populations of one tree each, Red Dadap and Sengon each there are two trees, For sampling, one tree each was taken in each type of *Fabaceae* and repeated sampling five times in each population sample.

5) Laboratory Analysis

The observation of stomatal type, stomatal number, and stomatal density on *Fabaceae* leaf types was conducted in the laboratory using a microscope with 400x magnification and a field of view area of 0.025 mm². Stomatal counting is done by counting the number of stomata visible in the observed area or a certain number of leaves. For example, count the number of stomata per unit of leaf area, such as mm² or cm². If different types of leaves or leaf parts are used, perform separate counts for each group. To quantify stomatal density, divide the number of stomata counted previously by the leaf area observed. For example, stomatal density can be expressed as the number of stomata per mm² or cm² of leaf. Stomatal density analysis is calculated using the formula below:

$$\text{Average stomata} = \frac{Sa_1 + Sa_2 + Sa_3 + \dots + Sa_n}{n}$$

$$\text{Somata density } (Ka_n) = \frac{\text{average } Sa}{FV}$$

Description:

Fv : Field View (mm²)

Sa₁ : Number of stomata in 1st field of view

Sa₂ : Number of stomata in 2nd field of view

Sa₃ : Number of stomata in 3rd field of view

Sa_n : Number of stomata in the field of view to (n)

6) Data Analysis

The data analysis method uses correlation analysis where the correlation analysis used is the correlation of persons (Crocker & Seber, 1980) to get the correlation value of Person (r) using the formula with the following equation (Stein & Corsten, 1991):

$$r_n = \frac{N \sum X.Y - \sum X \sum Y}{\sqrt{\{N \sum X^2 - (\sum X)^2\} \{N \sum Y^2 - (\sum Y)^2\}}}$$

Description:

R_{xy}: Person Product Moment Correlation Coefficient

x : Value of Independent Variabel

y : The Value of the Independent Variable

N : Number of Samples

Analysis was used with the SmartStatXL v.3.6.5.5 application to observe the effect of light intensity and air humidity on the number of stomata and stomatal density on the leaves of *Fabaceae* trees.

3. Result

1) Morphological characteristics of stomata on four *Fabaceae* species in the urban forest of Maros Regency

Fabaceae trees include plant species that are commonly found in the Urban Forest area and also become road protection trees. Based on observations in the research location of the Maros Regency urban forest, there are 4 types of *Fabaceae* plants consisting of Acacia (*Acacia auriculiformis*), Red Dadap (*Erythrina cristagalli* L), Sengon (*Paraserianthes falcataria* L), and Trembesi (*Samanea saman* (Jacq.) Merr.). The results of observations in the microscope to see the type of leaf stomata in four species of *Fabaceae* (*Acacia auriculiformis*, *Erythrina cristagalli* L, *Paraserianthes falcataria* L, *Samanea saman* (Jacq.) Merr.), in the Maros Regency Urban Forest can be seen in Figure 2. In Figure 2, it can be seen that the leaves of Acacia type have *diasitic* type of stomata, for red dadap Merah type has parasitic type of stomata, trembesi has *anomocytic* type of stomata, and sengon type has *cyclositic* type of stomata. Stomata in Acacia *auriculiformis* (Akasia) species with *diasitic* type can be seen in (Figure 1.A), where in the stomata there is a closing cell and surrounded by two neighbouring cells with a perpendicular location to the axis of the closing cell. In Figure 1. B stomata on the type of *Erythrina cristagalli* L (Dadap Merah) with parasitic stomata type, with the characteristics of the stomata, there is one closing cell accompanied by one or more neighboring cells with unequal size. The type of stomata in the *Samanea saman* (Jacq.) Merr.) (Trembesi) leaf type is *anomositik*, as shown in Figure 1. C can see the stomata have a cover cell surrounded by cells that resemble epidermal cells both in shape and size. The *Paraserianthes falcataria* L (Sengon) species with its stomatal type is the cyclomatic type, which has four or more neighboring cells surrounding the stomata in a circular pattern

(Figure 1.D). Although the four species observed are plant species from the same family, *Fabaceae*, they have different types of stomata in each species as obtained from the results of this study; this is because the habitat and water content of the

environment or place of growth are different (Prastika et al., 2023). Internal factors, namely the genetics of each species, cause differences in the type and number of stomata in each species. (Latifa et al., 2022).




Figure 2. Microscopic Viewing of *Fabaceae* Tree Stomata with 400 times magnification and a field of view area of 0.025 mm²

2) Number and Density of Stomata in four *Fabaceae* Tree Species in the Urban Forest of Maros Regency

The results of the analysis of the number and density of stomata in *Fabaceae* tree species (*Acacia*

auriculiformis, *Erythrina cristagalli* L, *Paraserianthes falcataria* L, *Samanea saman* (Jacq.) Merr.)) in the urban forest of Maros Regency are shown in Table 2.

Table 2. Number and Density of Stomata in four *Fabaceae* Tree Species in the Urban Forest of Maros Regency

| No | Tree Type of <i>Fabaceae</i> | Number of ES | Number of WS | Number of ES & WS | Density ES (mm ²) | Density WS (mm ²) | Number of stomata density ES & WS |
|--------------|--|----------------|----------------|-------------------|-------------------------------|-------------------------------|-----------------------------------|
| 1 | <i>Acacia auriculiformis</i> (Akasia) | 6.913 | 6.547 | 13.460 | 45.730,67 | 43.309,51 | 89.040,18 |
| 2 | <i>Erythrina cristagalli</i> L (Dadap Merah) | 414 | 396 | 810 | 2.738,68 | 2.619,60 | 5.358,28 |
| 3 | <i>Paraserianthes falcataria</i> L (Sengon) | 96.628 | 93.773 | 190.401 | 639.210,64 | 620.324,33 | 1.259.534,97 |
| 4 | <i>Samanea saman</i> (Jacq.) Merr.) (Trembesi) | 46.147 | 41.846 | 87.993 | 305.270,24 | 276.818,40 | 582.088,64 |
| Total | | 150.102 | 142.562 | 292.664 | 992.950,23 | 943.071,84 | 1.936.022,07 |

Description: ES= Eastern Stomata, WS= West Stomata

Of the four *Fabaceae* tree species shown in table 2, the Sengon species has the highest number of stomata (190,401) and stomatal density (1,259,534.97), while the Dadap Merah species has the lowest number of stomata (810) and stomatal density (5,358.28). Table 1 also shows that the number and density of Western stomata are lower than the number and density of Eastern stomata. The results showed that the higher the number of

stomata in plants, the higher the stomatal density. The density and number of stomata were influenced by environmental conditions, where the higher the light intensity, the higher the density (Budiono et al., 2016). In line with the research results (Meriko & Abizar, 2017) which state that the number of stomata affects the density of stomata, namely, if the number of stomata is large, the stomatal density is also higher. However, light intensity that is too high

is also not good for plant growth, so stomata need optimum light intensity to maximize stomatal work. In the morning, the light intensity is low, which optimizes the work of the stomata, compared to the afternoon, which has a high light intensity, resulting in excessive water evaporation.

3) Relationship between the number and density of stomata and environmental factors in the leaves of *Fabaceae* trees in the Maros Regency Urban Forest

Analysis of the relationship between the number and density of stomata to environmental factors on *fabaceae* tree leaves in the urban forest of maros regency is shown in table 3.

The correlation analysis of light intensity with the number of stomata and stomatal density showed (table 3) that only External Light Intensity of the Stand (ELIS) had a significant effect at the 0.05 (5%) level. In contrast, the Stand Light Intensity (SLI)

does not significantly affect the number of stomata and stomatal density. The results of this study indicate that any increase in External Light Intensity of the Stand (ELIS) will have a significant effect at the level of 0.05 (5%) on the ability of the number and density of Eastern stomata to absorb light by (-0.74), meaning that any increase in External Light Intensity of the Stand (ELIS) will reduce the ability of the number and density of Eastern stomata to absorb light intensity by -74%. The measurement results of the air humidity factor, both standing humidity (SH) and External Humidity of the Stand (EHS) in the correlation results, did not significantly affect the number of stomata and stomatal density. The number and density of stomata in each type of *Fabaceae* tree are different, where the highest number of stomata and stomatal density level is found in the type of session, while the number of stomata and the lowest stomatal density level is in the type of Dadap Merah

Table 3. Correlation Analysis of Light Intensity and Air Humidity on the Number and Density of Stomata in *Fabaceae*.

| Correlation Matrix Table (Pearson R) | | | | |
|--------------------------------------|--------------|------------|--------------|------------|
| | Number of ES | Density ST | Number of WS | Density WS |
| SLI | -0.26 | -0.26 | -0.22 | -0.22 |
| ELIS | -0.74 * | -0.74 * | -0.73 * | -0.73 * |
| SH | -0.09 | -0.09 | -0.09 | -0.09 |
| KLT | -0.08 | -0.08 | -0.09 | -0.09 |

Description: SLI= Stand Light Intensity, ELIS= External Light Intensity of the Stand, SH= Stand Humidity, EHS= External Humidity of the Stand, ES= Eastern Stomata, WS= West Stomata.

*) Significant at the real level of 0.05, The number on the body of the table is the value of the correlation coefficient.

The results of this study were supported by research (Carranza-Ramírez et al., 2025), which states that each plant cell has a different sensitivity and will affect the response in plants even given the same light intensity. The resulting research is also supported by research results (Juairiah, 2014), which state that differences in the number of stomata and stomatal density are influenced by internal factors, namely genetic differences, and external factors, namely the growing environment. (Rachmawati & Pujawati, 2020) stated that stomatal density is a genetic trait (heredity). This is supported by the results of research (Meriko & Abizar, 2017), which states that the level of stomatal density in each type of plant is different and influenced by the environment, such as water availability, temperature, CO₂ concentration, and light intensity. One of them is that the higher the light intensity, the higher the density of stomata on the leaf surface will also increase.

4. Discussion

The stomatal characteristics of the four *Fabaceae* species vary in terms of stomatal type, which differs among each species. This variation is influenced by environmental and genetic factors. Environmental factors affecting stomatal type include light intensity and air humidity. Although all six plant species

belong to the same family, they exhibit different stomatal types due to differences in their habitats and water availability in their growing environments (Prastika et al., 2023). Internal factors, such as the genetic makeup of each species, also play a significant role in determining the type and number of stomata (Latifa et al., 2022).

The stomatal type of the Acacia leaf is *diacytic*, characterized by two subsidiary cells surrounding the guard cells, positioned perpendicular to the long axis of the guard cells. Banyan and Capilong plants exhibit *paracytic* stomata, while palm has *mesoperigenous* stomata, and Acacia has *diacytic* stomata (Papuangan et al., 2014). The Red Dadap (*Erythrina*) species has *paracytic* stomata, where the long axis of the guard cells is parallel to the subsidiary cells, while the Trembesi (Rain Tree) species has *anomocytic* stomata, where the number of subsidiary cells surrounding the guard cells is indeterminate and indistinguishable from other epidermal cells.

Three types of stomata were observed in this study: the Rain Tree (Ki Hujan) leaf has *anomocytic* stomata, the Butterfly Tree (Kupu-kupu) leaf has *anisocytic* stomata, and the Red Dadap (Dadap Merah) has *paracytic* stomata (Jaya et al., 2014). The Sengon (*Albizia*) species exhibits *cyclocytic* stomata, where four or more subsidiary cells form a

ring-like structure around the stomata. *Cyclocytic* stomata are characterized by four or more subsidiary cells arranged in a circular pattern surrounding the stomata (Evert, 2006).

The results of the study on the number and density of stomata in *Fabaceae* plants indicate that the number and density of stomata on the western side are lower compared to those on the eastern side. The higher the light intensity received by the trees, the greater the number and density of stomata. Stomatal density and number are influenced by environmental conditions, where higher light intensity leads to higher stomatal density (Budiono et al., 2016). Additionally, this is supported by research from (Meriko & Abizar, 2017), which states that the number of stomata affects stomatal density, meaning that a higher number of stomata results in greater stomatal density. However, excessively high light intensity is also not beneficial for plant growth, as stomata require optimal light intensity to function efficiently. In the morning, light intensity is not too high, allowing stomata to work more optimally compared to midday, when light intensity is high, leading to excessive water evaporation. Genetic factors and the differing environmental conditions of the plant's habitat also influence stomatal characteristics. Furthermore, light intensity, water availability, temperature, and CO₂ concentration are additional factors that affect the number and density of stomata (Juma'ani & Munawwaroh, 2017).

The number and density of stomata vary among each *Fabaceae* tree species, with the highest stomatal count and density found in the Sengon (*Albizia*) species, while the lowest stomatal count and density are observed in the Dadap Merah (*Erythrina*) species. These findings are supported by research from (Salisbury & Ross, 1995), which states that each cell in plants has varying sensitivities, influencing the plant's response even when exposed to the same light intensity. The results are further corroborated by a study from (Juairiah, 2014), which explains that internal factors, such as genetic variation, and external factors, such as the environmental conditions of the growing site influence differences in stomatal nu. Rachmawati and Pujawati (2020) state that stomatal density is a genetic trait (inherited). This is supported by research from (Meriko & Abizar, 2017), which indicates that stomatal density varies among plant species and is influenced by environmental factors such as water availability, temperature, CO₂ concentration, and light intensity. For instance, higher light intensity leads to an increase in stomatal density on the leaf surface.

The relationship between stomatal number, stomatal density, and environmental factors—specifically light intensity and air humidity—shows that only External Light Intensity has a significant influence. Each increase in External Light Intensity affects the ability of the stomata on the eastern and western sides to absorb light. As External Light Intensity increases, the stomata tend to close to prevent excessive water evaporation, thereby reducing light absorption by the stomata. This is supported by research from Sumadji (2020), which states that stomatal closure is an adaptation mechanism to drought stress, reducing

transpiration rates and preventing water loss during periods of high light intensity and temperature. Additionally, research by Sudomo (2009) indicates that both extremely low and extremely high light intensities negatively impact plant growth by slowing it down, as they affect the functioning of stomatal cells in the leaves. Each tree species has an optimal range of light intensity, within which stomatal function can operate efficiently. Sudomo (2009) further emphasizes that optimal light intensity is crucial for maximizing plant growth. However, Light Intensity, Stand Humidity, and External Humidity do not have a significant effect on stomatal number and density. Future research should explore the effects of varying CO₂ concentrations and soil moisture on stomatal behavior, as these factors were not addressed in this study. Additionally, long-term monitoring of stomatal adaptations to seasonal changes and extreme climate conditions could provide deeper insights into plant resilience. Genetic analysis of stomatal traits and comparative studies across different ecosystems would further enhance our understanding of stomatal adaptations in *Fabaceae* species

5. Conclusion

This study analysed the relationship of stomatal characteristics in *Fabaceae* tree leaves with environmental factors, specifically light intensity and air humidity in the Maros City Forest. The results showed that the type, number and density of stomata varied among *Fabaceae* species, reflecting adaptation to specific environmental conditions. External light intensity (ELIS) had a significant effect on stomatal number and density with a negative correlation at the 5% significance level. In contrast, light intensity within the stand (SLI) had no significant effect because the trees provided shading that stabilised the light intensity. Air humidity inside and outside the stand had no significant impact on stomata as the variation was small in the urban forest environment. Other factors, such as plant genetics and carbon dioxide concentration, are more dominant in influencing stomatal characteristics.

This study has limitations, such as not considering seasonal variability as well as other factors such as carbon dioxide concentration, soil quality, and air temperature. Addressing these variables in future research could provide a more comprehensive understanding of the environmental drivers influencing stomatal behaviour. Further research is recommended to explore the genetic mechanisms underlying stomatal differentiation and to expand the study to other plant families within the Maros Regency urban forest. Such investigations would enhance our knowledge of plant adaptive strategies and support the development of effective conservation practices. Moreover, integrating physiological measurements, such as transpiration rates and photosynthetic efficiency, could provide a more holistic understanding of how stomatal behaviour influences plant ecology.

6. Author Contributions

The first author (RY) contributes as a compiler and data analyzer based on the research results. In contrast, the second author (HS) contributes methods to help interpret data and research results, and the third (N) and fourth authors (IND) contribute to help prepare the article.

7. Competing Interests

All authors declare no conflicts of interest regarding this publication.

8. Acknowledgements

Thanks to the Indonesian Research and Technology Agency (BRIN) for providing funding for thesis research through the BARISTA program, the Hasanuddin University Faculty of Agriculture Integrated Laboratory for its facility support, and to the Supervisors from BRIN and the Forestry Study Program, Faculty of Agriculture and Forestry, Maros Muslim University who have spent much time for the preparation of this research

9. References

- Budiono, R., Sugiarti, D., Nurzaman, M., Setiawati, T., Spriatun, T. & Mutaqien, A. Z. (2016). Kerapatan Stomata dan Kadar Klorofil Tumbuhan *Clausena excavata* Berdasarkan Perbedaan Intensitas Cahaya. Seminar Nasional Pendidikan Dan Saintek UNPAD: FMIPA Biologi, 2016, 61–65. <https://proceedings.ums.ac.id/snpbs/article/view/363>
- Cambaba, S. & Kasi, P. D. (2022). Karakteristik Stomata Daun Pucuk Merah (*Syzygium oleana*) Berdasarkan Waktu Pengambilan Sampl Yang Berbeda. *Cokroaminoto Journal of Biological Science*, 4(1), 19-25. Retrieved from <https://science.e-journal.my.id/cjbs/article/view/95>
- Carranza-Ramírez, J. E., Borda, A. M. & Moreno-Fonseca, L. P. (2025). LED light modifies plant architecture, physiological parameters and cannabinoid content in three varieties of *Cannabis sativa* L. *South African Journal of Botany*, 176, 231–240. <https://doi.org/10.1016/j.sajb.2024.11.023>
- Casson S, Gray JE. Influence of environmental factors on stomatal development. *New Phytol.* 2008;178(1):9-23. <https://doi.org/10.1111/j.1469-8137.2007.02351.x>
- Chauhan, J., Prathibha, M. D., Singh, P., Choyal, P., Mishra, U. N., Saha, D., Kumar, R., Anuragi, H., Pandey, S., Bose, B., Mehta, B., Dey, P., Dwivedi, K. K., Gupta, N. K. & Singhal, R. K. (2023). Plant photosynthesis under abiotic stresses: Damages, adaptive, and signaling mechanisms. In *Plant Stress* (Vol. 10). Elsevier B.V. <https://doi.org/10.1016/j.stress.2023.100296>
- Clark, J. W., Harris, B. J., Hetherington, A. J., Hurtado-Castano, N., Brench, R. A., Casson, S., Williams, T. A., Gray, J. E. & Hetherington, A. M. (2022). The origin and evolution of stomata. In *Current Biology* (Vol. 32, Issue 11, pp. R539–R553). Cell Press. <https://doi.org/10.1016/j.cub.2022.04.040>
- Crocker, D. C. & Seber, G. A. F. (1980). Linear Regression Analysis. In *Technometrics* (Vol. 22, Issue 1). <https://doi.org/10.2307/1268395>
- Demars, B. O. L., Schneider, S. C., Thiemer, K., Dörsch, P., Pulg, U., Stranzl, S., Velle, G. & Pathak, D. (2024). Light and temperature controls of aquatic plant photosynthesis downstream of a hydropower plant and the effect of plant removal. *Science of the Total Environment*, 912. <https://doi.org/10.1016/j.scitotenv.2023.169201>
- Djahangard, M., Buckwitz, S., Hioki, C., He, J., Zhou, Z., Zhang, H. & Yousefpour, R. (2024). A global comparative analysis of local recreation behaviors and values in peri-urban forests. *Environmental and Sustainability Indicators*, 24. <https://doi.org/10.1016/j.indic.2024.100525>
- Escobedo, F. J., Kroeger, T., & Wagner, J. E. (2011). Urban forests and pollution mitigation: Analyzing ecosystem services and disservices. *Environmental Pollution*, 159(8-9), 2078–2087. <https://doi.org/10.1016/j.envpol.2011.01.010>
- Evert, R.F. (2006). *Esau's Plant Anatomy (Meristems, Cell and Tissues of the PlantBody: Their Structure, Function and Development*. 3rd. A John Wiley & Sons. Inc. Publication. New Jersey
- Firmansyah, A., Rahmawati, T., Hardiyanti, W., Meylia, S. A., Yulastri, W., Asidqi, A. & Rojulun Afif, V. (2022). Tumbuhan Terancam Punah di Hutan Kota Ranggawulung Subang. 77.
- Franks PJ, Farquhar GD. The mechanical diversity of stomata and its significance in gas-exchange control. *Plant Physiol.* 2007 Jan;143(1):78-87. <https://doi.org/10.1104/pp.106.089367>
- Handayani, N. P., Rahmadania, A. P., Annisa, Z. D., Haryanti, A., Purwaningrum, I. F., Sudaryoko, D. A. P., Febriana, B. W. S., Balerina, A. M. & Rahayu, T. D. (2024). Upaya Pengurangan Polusi Udara di Lingkungan Universitas Negeri Semarang dengan Penanaman Pohon Jurnal Majemuk, 3.2(2), 256–268. <http://jurnalilmiah.org/journal/index.php/majemuk>
- Haworth, M., Marino, G., Atzori, G., Fabbri, A., Daccache, A., Killi, D., Carli, A., Montesano, V., Conte, A., Balestrini, R., & Centritto, M. (2023). Plant Physiological Analysis to Overcome Limitations to Plant Phenotyping. *Plants*, 12(23), 4015. <https://doi.org/10.3390/plants12234015>
- Hetherington AM, Woodward FI. The role of stomata in sensing and driving environmental

- change. *Nature*. 2003 Aug 21;424(6951):901-8. <https://doi.org/10.1038/nature01843>.
- Jaya, Abrar Brata, Elis Tambarua, A. Ilham Latunrab, and Muhtadin Asnady Salamc. (2014). "Perbandingan Karakteristik Stomata Daun Pohon Leguminosae Di Hutan Kota Universitas Hasanuddin Dan Di Jalan Tamalate Makassar." oai: repository.unhas.ac.id:123456789/13229
- Juairiah, L. (2014). Studi Karakteristik Stomata Beberapa Jenis Tanaman Revegetasi Di Lahan Pascapenambangan Timah Di Bangka. *Widyariset*, 17(2), 213-217. Juma'Ani, Juma'Ani, and Anita Munawwaroh. (2017). "Analisis Karakteristik Stomata Pada Daun Tanaman Bambu Rejeki (*Dracaena Reflexa*) Sebagai Tanaman Hias Penyerap Polusi Di Kawasan Kota Malang." *Edubiotik*, vol. 2, no. 02, pp. 7-12, <https://doi.org/10.33503/ebio.v2i02.125>.
- Juslina. (2022). Analisis kandungan klorofil enam jenis pohon di kampus universitas hasanuddin tamalanrea. <http://repository.unhas.ac.id:443/id/eprint/18778>
- Latifa, R., Nurrohman, E. & Hadi, S. (2022). Stomata leaves characteristics of sapindaceae family in malabar forest, Malang city. *Bioscience*, 6(2), 73. <https://doi.org/10.24036/0202262118189-0-00>
- Lawson T, Blatt MR. Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiol*. 2014 Apr;164(4):1556-70. <https://doi.org/10.1104/pp.114.237107>.
- Merdekawati, R. P. (2015). Jumlah dan Ukuran Stomata pada Daun Glodokan (*Polyalthia longifolia*) di Jalan Tun Abdul Razak dan di Area Kampus UIN Alauddin Makassar. [Skripsi]. Makassar: Fakultas Sains Dan Teknologi Uin Alauddin Makassar.
- Meriko, L. & Abizar. (2017). Struktur Stomata Daun Beberapa Tumbuhan Kantong Semar (*Nepenthes* spp.) [Structure of Leaves Stomata on Some Pitcher Plants (*Nepenthes* spp.)]. *Jurnal Ilmu-Ilmu Hayati*, 16(3), 325-330. <https://doi.org/10.14203/beritabiologi.v16i3.2398>
- Nowak, D. J., Hirabayashi, S., Bodine, A., & Greenfield, E. (2014). Tree and forest effects on air quality and human health in the United States. *Environmental Pollution*, 193, 119-129. <https://doi.org/10.1016/j.envpol.2014.05.028>
- Ordóñez, C., & Duinker, P. N. (2013). An analysis of urban forest management plans in Canada: Implications for urban forest management. *Landscape and Urban Planning*, 116, 36-47. <https://doi.org/10.1016/j.landurbplan.2013.04.007>
- Papuan. N., Nurhasanah., M. D. (2014). Jumlah Dan Distribusi Stomata Pada Tanaman Penghijauan Di Kota Ternate. 3(September). <https://doi.org/https://doi.org/10.33387/bioedu.v2i1.62>
- Prastika, D., Sarjani, T. M., Mahyuni, S. R., Hariani, I., Ramadhan, D. A., Rezeki, S., Tiara, R., Hendrik, E., Aulia, R. & Amalia, T. (2023). Identifikasi Tipe Stomata Anggota Suku Myrtaceae di Kota Langsa. *Jurnal Sains Dan Edukasi Sains*, 6(1), 20-27. <https://doi.org/10.24246/juses.v6i1p20-27>
- Rachmawati, R. N. & Pujawati, E. D. (2020). Karakteristik Stomata Nyawai (*Ficus variegata* Blume) dari 3 Sumber Benih Asal Kalimantan di KHDTK Riam Kiwa Desa Lobang Baru The Characteristics of Stomata Nyawai (*Ficus variegata* Blume) From 3 Sources of Origin Seeds Kalimantan in KHDTK Riam Kiwa Lobang. *Jurnal Sylva Scientae*, 3(6), 1078-1085. <https://doi.org/10.20527/jss.v3i6.4725>
- Raharjo, H. P., Haryanti, S., & Hastuti, R. B. (2015). Pengaruh Tingkat Kepadatan Lalu Lintas dan Waktu Pengamatan yang Berbeda Terhadap Ukuran dan Jumlah Stomata Daun Glodokan (*Polyalthia longifolia*. Sonn). *Jurnal Akademika Biologi*, 4(1), 73-84. Retrieved from <https://ejournal3.undip.ac.id/index.php/biologi/article/view/19402>
- Salisbury FB, Ross CW. (1995). *Plant physiology*. 4th ed. Wadsworth Publishing Co, Belmont, California.
- Sinay, H. & Lesilolo, M. H. (2020). Karakterisasi Stomata Daun Pegagan (*Centella asiatica* (L.) Urban.) pada Ketinggian Wilayah Berbeda di Pulau Ambon. *Science Map Journal*, 2(1), 1-7. <https://doi.org/10.30598/jmsvol2issue1pp1-7>
- Stein, A. & Corsten, L. C. A. (1991). Universal Kriging and Cokriging as a Regression Procedure. *Biometrics*, 47(2), 575. <https://doi.org/10.2307/2532147>
- Sudomo, A. (2009). Pengaruh naungan terhadap pertumbuhan dan mutu bibit manglid (*Manglieta glauca* BI). *Tekno hutan tanaman*, 2(2), 59-66.
- Sulistiyowati, Hari, Emitria Rahmawati, and Retno Wimbaningrum. 2021. "Spatial Distribution Patterns of Lantana camara L. Population as Invasive Alien Species In Pringtali Savana Bandealet Resort Meru Betiri National Park." *Jurnal ILMU DASAR* 22(1): 19.
- Sumadji, Angga Rahabistara. (2020). "Kerapatan Stomata Dan Kaitannya Terhadap Kekeringan Pada Tanaman Padi Varietas IR64." *Widya Warta Jurnal Ilmiah Tahun XLIV*(1): 43-54. <http://portal.widyamandala.ac.id/jurnal/index.php/warta/article/view/907>
- Suwandi, R. (2021). Fungsi dan Manfaat Ruang Terbuka Hijau Di Kelurahan Kampung Bandar Kecamatan Senapelan Kota Pekanbaru.

Universitas Islam Riau.
http://repository.uir.ac.id/id/eprint/18805
World Health Organization (WHO). (2016). Urban
green spaces and health: A review of

evidence. Copenhagen: WHO Regional Office
for Europe.

How to cite this article:

Yahya, R. Y., Sukri, H., Nirawati, & Djarot, I. N. D. (2025). Responses of Stomatal Characteristics to Environmental Factors in Fabaceae Trees of the Urban Forest in Maros Regency. *Jurnal Wasian*, 12(01), 01-09. <https://doi.org/10.62142/xx0wes03>



[Endnote/Zotero/Mendeley\(RIS\)](#)



[BibTex](#)